

## **Infection Control in Interscholastic Athletic Training Rooms Post COVID-19**

### **Problem**

COVID-19 has caused our entire population to re-evaluate how we react and relate to each other's physical space, and how this impacts our collective hygiene and health in order to minimize the transmission of fungal, viral or bacterial disease. We are now more conscious of our proximity with one another, and who or what we do, or do not, touch.

Personal hand hygiene, personal distancing, and the clean dis-infected status of objects or surfaces we commonly touch are now everyday considerations. As individuals, we expect the professionals and clinical sites providing us with our health care to maintain the highest standards of cleanliness and infection control possible to minimize our risk. These standards must also be maintained at the location where athletes commonly seek assistance on a daily basis during the school year, the athletic training room. The ability of the professional staff to meet these standards is often severely challenged based upon the volume of athletes they need to assist in a specific time period, the physical location and size of the athletic training room and school cleaning standards.

### **Background**

Interscholastic athletic injuries are usually musculoskeletal in nature and the health care issue most commonly presented by middle and high school athletes to athletic training rooms.<sup>1-3</sup> However, athletic trainers also commonly work with athletes who have non-musculoskeletal related illnesses and infections on an acute and sub-acute basis. These issues typically have revolved around skin infections, include multidrug-resistant organisms, such as vancomycin-resistant enterococcus (VRE) and methicillin-resistant staphylococcus aureus (MRSA), from close contact associated with some sports and ill-defined cleaning standards.<sup>4-9</sup>

Though commonly thought of as being associated only with skin lesions and infections, MRSA can be a potential source of bacterial wound infections, pneumonia, and urinary tract infections. Any of these MRSA based infections may have a high morbidity that can require hospitalization and antibiotic therapy. Athletes with other types of viral and fungal infections may also be evaluated and triaged by athletic trainers and then referred on to other health care professionals, such as the school nurse, team physician, school physician, or athletes' personal physicians. The 2020 development, presence and prevalence of COVID-19 throughout the country and the significant health care concerns associated with COVID-19 have heightened the need for vigilance in maintaining optimal infection control standards in any athletic health care area.

The typical high school athletic training room is usually a small space that is a shared environment. This shared environment is most often the primary athletic health care clinical environment for the injured or ill athlete. Unfortunately, this area is all too often also commonly used for the storage of supplies, coolers, ice machines, and other equipment. This shared environment and tight space often facilitates the close contact between surfaces and objects, multiple individual athletes, and groups of athletes all seeking the attention of the athletic trainer for their specific needs during a short time period as they get ready for practice or games. As a shared environment, the athletic training room can act as a source for the spread of infection especially in the presence of poor hygiene and contamination.<sup>10</sup> Due to limitations of space and funding, most high school athletic

training rooms have difficulty meeting standards suggested for optimal provision of athletic health care needs and infection control.<sup>11-14</sup>

Studies have demonstrated that the establishment and adherence to appropriate infection control policies and procedures, the education of staff, coaches, parent and students, and the utilization of the appropriate products can specifically reduce bacterial and viral burdens (including multi-drug resistant organisms).<sup>4,6,7,10</sup> Greater attention and detail to improve infection control and facility standards are necessary to assure the health of all athletes and minimize the transmission of communicable disease in this COVID and Post COVID-19 environment.

## **Solution**

Establish and maintain partnership between the school and Brigham Health Care Department of Orthopedics and Sports Medicine Service for the provision of the Athletic Health Care needs of the school's student athletes.

Brigham Health Care is required by licensure to adhere to the highest infection control standards for all services provided. Through partnership with Brigham Health Care and the contractual provision of athletic health care services, the following would be initiated:

1. Identifying and assigning a team physician and certified athletic trainer responsible for oversight of all athletic health care needs of the school's athletes in collaboration with other school health providers, including the school physician, school nurse, and other school health providers associated with the school.
2. Collaborative development of policies and procedures that meet commonly accepted medical and professional standards for infection control in the athletic training room, such as:
  - a. Infection control and cleaning standards within, and in the immediate adjacent area to, the training room;
  - b. Cleaning and storage standards for athletic health care equipment and supplies;
  - c. Review and recommendations for physical space associated with athletic training room to enhance ability to meet infection control standards, policies and procedures based upon BOC Facility Principles and NATA Appropriate Medical Care Standards; and
  - d. Development of an educational program for athletic department staff, coaches, parents and athletes regarding need for appropriate hygiene and infection control standards to minimize exposure to athletes.

## **References**

1. Collins CL, Comstock RD. Epidemiological features of high school baseball injuries in the United States, 2005-2007. *Pediatrics* 2008; 121(6):1181-1187.
2. Rechel JA, Yard EE, Comstock RD. *J Athletic Training* 2008; 43(2):197-204.
3. Powell JW, Barber-Foss KD. Injury patterns in selected high school sports: A review of the 1995-1997 seasons. *J Athletic Training* 1997; 34(3):277-284.
4. Ferrara MS, Courson R, Paulson DS. Evaluation of persistent antimicrobial effects of an antimicrobial formulation. *J Athletic Training* 2011; 46(6):629-633.
5. Zinder SM, Basler RW, Foley J et al: National athletic trainers' association position statement: Skin diseases. *J Athletic Training* 2010; 45(4):411-428.

6. Young LM, Abram V, Markey ER et al. Recommendations for best disinfectant practices to reduce the spread of infection via wrestling mats. J Athletic Training 2017; 52(2):82-88.
7. Messina G, Burgassi S, Russo C et al. Is it possible to sanitize athletes' shoes? J Athletic Training 2015; 50(2):126-132.
8. Jimenez-Truque N, Saye EJ, Soper N et al. Association between contact sports colonization with staphylococcus aureus in a prospective cohort of collegiate athletes Sports Med 2017; 47:1011-1019.
9. Kahanov L, Kim YK, Eberman et al. A staphylococcus aureus and community-associated methicillin resistant staphylococcus aureus (CA-MRSA) in and around therapeutic Whirlpools in athletic training rooms. J Athletic Training 2015; 50(4):432-437.
10. Labelle MW, Knapik DM, Arbogast JW et al. Infection risk reduction program on pathogens in high school and collegiate athletic training rooms. Sports Health 2020; 12(1):51-57.
11. [https://www.nata.org/sites/default/files/nata\\_appropriate\\_medical\\_care\\_standards.pdf](https://www.nata.org/sites/default/files/nata_appropriate_medical_care_standards.pdf)
12. <https://www.nata.org/sites/default/files/communicableinfectiousdiseasessecondaryschoolsports.pdf>
13. [https://bocatc.org/system/document\\_versions/versions/42/original/boc-facility-principles-20170615.pdf?1497543426](https://bocatc.org/system/document_versions/versions/42/original/boc-facility-principles-20170615.pdf?1497543426)
14. Courson R, Goldberg M, Adams KG et al. Inter-association consensus statement on best practices for sports medicine management for secondary schools and colleges. J Athletic Training 2014; 49(1):128-137.

Developed By:

Jim Zachazewski, ATC, PT, DPT

Athletic Trainer

Department of Orthopedic Surgery – Sports Medicine

Brigham and Women's Hospital

Boston, MA

[jzachazewski@partners.org](mailto:jzachazewski@partners.org)

Michael Belanger, ATC, PT

Clinical Supervisor of Athletic Training

Department of Orthopedic Surgery – Sports Medicine

Brigham and Women's Hospital

Boston, MA

[mjbelanger@bwh.harvard.edu](mailto:mjbelanger@bwh.harvard.edu)